MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 3, 2021/2022

BMS 1024 – MANAGERIAL STATISTICS

(All sections / Groups)

04 OCTOBER 2022
2.30 p.m. – 4.30 p.m.
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This question paper consists of **NINE (9) pages** excluding cover page with **3 Questions** only.

2. Answer all **THREE (3) questions**

3. Formula and Statistical tables are attached at the end of the question paper.

4. Students are allowed to use non-programmable scientific calculators with no restrictions.

Question 1 (25 Marks)

a) From an inventory of 20 cars being shipped to local automobile dealers, 5 have defective radios installed. Three cars are selected at random and tested. Let X be the random variable that denotes the number of cars having defective radios installed

i) Construct the probability distribution of X. **(8 marks)**

ii) Find the standard deviation of X. **(4 marks)**

b) According to the survey conducted by Rakuten Insight in 2021, 24 % of the Malaysian respondents ordered food on delivery apps once a week. Assume that 10 Malaysian respondents are randomly selected, find the probability that 3 of the selected respondents ordered food on delivery apps once a week. (Round to four decimal places). **(3 marks)**

c) The number of users of dryer in a laundry self-service shop is Poisson distributed. The mean number of users per 1-hour interval is 1.5. Find the probability that there will be less than 2 users in the next three hours. (Round to four decimal places). **(5 marks)**

d) A packaging bag are manufactured so that the breaking strength of the bag is normally distributed with a mean of 6 pound per square inch and a standard deviation of 2.2 pound per square inch. What proportion of the bag produced have a breaking strength of at least 5.2 pounds per square inch? Sketch your normal distribution. **(5 marks)**

Question 2 (55 Marks)

a) Suppose that in Malaysia, the monthly income of insurance agents is normally distributed. A random sample of 10 insurance agents was asked to report their monthly income (in RM100) and listed as below:

47	46	42	48	50
49	45	43	42	43

i) Calculate the mean and standard deviation by using formula. **(5 marks)**

ii) Construct a 95% confidence interval estimate of the population mean. Express your answer in 2 decimal places. **(5 marks)**

iii) At 90% confidence level, how large a sample should be selected if they want the estimate to be within RM0.50 of the population mean? Assume that the population standard deviation is known to be RM3. **(5 marks)**

Continued...

b) According to the Centers for Disease Control and Prevention (CDC), a height of ten years old girls is normally distributed with a mean of 60 inches. A researcher claims that the height of girls who live in Malaysia will have a lower height compared to the population mean. To test the researcher's claim, a random sample of 40 girls were examined. The mean height in this sample was 55.5 inches. Would you conclude that the researcher's claim is true at a 5% significance level? Assume that the population standard deviation is known to be 15 inches.

(10marks)

c) On average, the academic performance of female students measured by CGPA in a university are higher compared to male students. To test that hypothesis, a random sample of 20 students was chosen. Their CGPA result were recorded in the following table.

	Female	Male
Sample size	10	10
Mean, $\bar{X}$	2.982	2.972
Standard deviation, s	0.360	0.458

Is there enough evidence at the 5% significance level to infer that female students perform better than male students? **(10 marks)**

d) The Excel output provided below is to study the effect of income (one unit of income = \$10,000) has on the happiness (where happiness is measured by scale of 1 to 10) using the simple linear regression model for a sample size of 12 peoples.

ANOVA

	df	*SS*	*MS*	*F*	*Significance F*
Regression	1	1124.315	1124.32	4.063492	0.1974922
Residual	10	376.1286	0.75832		
Total	11	1500.443			

	Coefficients	*Standard Error*	*t Stat*	*P-value*
Intercept	0.20427	0.764398	1.17289	0.361624
Income	0.71383	0.422704	0.79599	0.509506

i) Determine the dependent variable (Y) and independent variable (X) for the regression model. **(2 marks)**

ii) Determine the simple regression equation. **(3 marks)**

iii) Interpret the slope and intercept coefficients in ii). **(4 marks)**

iv) Calculate the coefficient of determination, R^2. Interpret the value. **(4 marks)**

v) Conclude at 5% significance level, is there a linear relationship between happiness and income? Use the p-value approach. **(7 marks)**

Continued...

Question 3 (20 Marks)

a) A research has been done to compare the sale prices for year 2020 and 2021. Create a simple index by using 2020 as the base year. **(7 marks)**

Commodity	2020	2021
A	30	20
B	70	50
C	90	110
D	100	120
E	130	100
F	70	50
G	50	80

b) The following table shows three type of goods and their prices for year 2020 and 2021.

Good	2020 Price	2020 Quantity	2021 Price	2021 Quantity
A	0.25	60	0.28	80
B	0.35	120	0.4	210
C	0.3	40	0.45	60

Determine and interpret the Laspeyres' Price Index, Paasche's Price Index and Fisher's Price Index for A, B and C by using 2020 as the base period.

(13 marks)

End of Page.

STATISTICAL FORMULA

A. DESCRIPTIVE STATISTICS

Mean $(\bar{x}) = \frac{\sum_{i=1}^{n} X_i}{n}$

Standard Deviation $(s) = \sqrt{\frac{\sum_{i=1}^{n} X_i^2}{n-1} - \frac{(\sum_{i=1}^{n} X_i)^2}{n(n-1)}}$

B. PROBABILITY

Poisson Probability Distribution

If X follows a Poisson Distribution , $P(\lambda)$ where $P(X = x) = \frac{e^{-\lambda}\lambda^x}{x!}$

then the mean = $E(X) = \lambda$ and variance = $VAR(X) = \lambda$

Binomial Probability Distribution

If X follows a Binomial Distribution B(n, p) where $P(X = x) = {}^nC_x p^x q^{n-x}$

then the mean = $E(X) = np$ and variance = $VAR(X) = npq$ where $q = 1 - p$

Normal Distribution

If X follows a Normal distribution, $N(\mu, \sigma)$ where $E(X) = \mu$ and $VAR(X) = \sigma^2$

then $Z = \frac{X - \mu}{\sigma}$

C. EXPECTATION AND VARIANCE OPERATORS

$E(X) = \sum[X \bullet P(X)]$

$VAR(X) = E(X^2) - [E(X)]^2$ where $E(X^2) = \sum[X^2 \bullet P(X)]$

If $E(X) = \mu$ then $E(cX) = c\mu$, $E(X_1 + X_2) = E(X_1) + E(X_2)$

If $VAR(X) = \sigma^2$ then $VAR(cX) = c^2\sigma^2$,

$VAR(X_1 + X_2) = VAR(X_1) + VAR(X_2) + 2\,COV(X_1, X_2)$

where $COV(X_1, X_2) = E(X_1X_2) - [E(X_1)\,E(X_2)]$

D. CONFIDENCE INTERVAL ESTIMATION AND SAMPLE SIZE DETERMINATION

(100 – α) % Confidence Interval for Population Mean (σ Known) $\mu = \bar{X} \pm Z_{\alpha/2}\left(\sigma/\sqrt{n}\right)$

(100 – α)% Confidence Interval for Population Mean (σ Unknown) $\mu = \bar{X} \pm t_{\alpha/2,n-1}\left(s/\sqrt{n}\right)$

(100 – α)% Confiden .: Interval for Population Proportion $p = \hat{p} \pm Z_{\alpha/2}\sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$

Sample Size Determination for Population Mean	$n = \frac{Z^2\sigma^2}{E^2}$
Sample Size Determination for Population Proportion	$n \geq \frac{Z^2\,\hat{p}(1-\hat{p})}{E^2}$
Where E = Limit of Error in Estimation	

E. HYPOTHESIS TESTING

One Sample Mean Test

Standard Deviation (σ) Known	Standard Deviation (σ) Not Known
$Z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$	$t = \frac{\bar{x} - \mu}{s/\sqrt{n}}$

One Sample Proportion Test

$$Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$$

Two Sample Mean Test

Standard Deviation (σ) Known	Standard Deviation (σ) Not Known
$z = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\sigma_1^2/n_1 + \sigma_2^2/n_2}}$	$t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{S_p^2\left(1/n_1 + 1/n_2\right)}}$ where $S_p^2 = \frac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{(n_1 + n_2 - 2)}$

Two Sample Proportion Test

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{p(1-p)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \quad \text{where } p = \frac{X_1 + X_2}{n_1 + n_2}$$

where X_1 and X_2 are the number of successes from each population

F. REGRESSION ANALYSIS

Simple Linear Regression

Population Model: $Y = \beta_0 + \beta_1 X_1 + \varepsilon$

Sample regression equation: $\hat{y} = b_0 + b_1 x_1$

Correlation Coefficient

$$r = \frac{\sum XY - \left[\frac{\sum X \sum Y}{n}\right]}{\sqrt{\left[\sum X^2 - \left(\left(\sum X\right)^2/n\right)\right]\left[\sum Y^2 - \left(\left(\sum Y\right)^2/n\right)\right]}} = \frac{COV(X,Y)}{\sigma_X \sigma_Y}$$

ANOVA Table for Regression			
Source	**Degrees of Freedom**	**Sum of Squares**	**Mean Squares**
Regression	*1*	SSR	MSR = SSR/1
Error/Residual	$n-2$	SSE	MSE = SSE/$(n-2)$
Total	$n-1$	SST	

Test Statistic for Significance of the Predictor Variable

$t_i = \dfrac{b_i}{S_{b_i}}$ and the critical value = $\pm t_{\alpha/2,(n-p-1)}$

Where p= number of predictor

G. INDEX NUMBERS

Simple Price Index $P = \dfrac{p_t}{p_0} \times 100$	**Laspeyres Quantity Index** $P = \dfrac{\sum p_0 q_t}{\sum p_0 q_0} \times 100$
Aggregate Price Index $P = \dfrac{\sum p_t}{\sum p_0}(100)$	**Paasche Quantity Index** $P = \dfrac{\sum p_t q_t}{\sum p_t q_0} \times 100$
Laspeyres Price Index $P = \dfrac{\sum p_t q_0}{\sum p_0 q_0} \times 100$	**Fisher's Ideal Price Index** $\sqrt{(Laspeyres\ \ Price\ Index)(Paa\ sche\ Price\ Index)}$
Paasche Price Index $P = \dfrac{\sum p_t q_t}{\sum p_0 q_t} \times 100$	**Value Index** $V = \dfrac{\sum p_t q_t}{\sum p_0 q_0} \times 100$

B. STATISTICAL TABLE

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

The function tabulated is $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-\frac{1}{2}t^2}\, dt$. $\Phi(x)$ is the probability that a random variable, normally distributed with zero mean and unit variance, will be less than or equal to x. When $x < 0$ use $\Phi(x) = 1 - \Phi(-x)$, as the normal distribution with zero mean and unit variance is symmetric about zero.



x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
0·00	0·5000	0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·97725
·01	·5040	·41	·6591	·81	·7910	·21	·8869	·61	·9463	·01	·97778
·02	·5080	·42	·6628	·82	·7939	·22	·8888	·62	·9474	·02	·97831
·03	·5120	·43	·6664	·83	·7967	·23	·8907	·63	·9484	·03	·97882
·04	·5160	·44	·6700	·84	·7995	·24	·8925	·64	·9495	·04	·97932
0·05	0·5199	0·45	0·6736	0·85	0·8023	1·25	0·8944	1·65	0·9505	2·05	0·97982
·06	·5239	·46	·6772	·86	·8051	·26	·8962	·66	·9515	·06	·98030
·07	·5279	·47	·6808	·87	·8078	·27	·8980	·67	·9525	·07	·98077
·08	·5319	·48	·6844	·88	·8106	·28	·8997	·68	·9535	·08	·98124
·09	·5359	·49	·6879	·89	·8133	·29	·9015	·69	·9545	·09	·98169
0·10	0·5398	0·50	0·6915	0·90	0·8159	1·30	0·9032	1·70	0·9554	2·10	0·98214
·11	·5438	·51	·6950	·91	·8186	·31	·9049	·71	·9564	·11	·98257
·12	·5478	·52	·6985	·92	·8212	·32	·9066	·72	·9573	·12	·98300
·13	·5517	·53	·7019	·93	·8238	·33	·9082	·73	·9582	·13	·98341
·14	·5557	·54	·7054	·94	·8264	·34	·9099	·74	·9591	·14	·98382
0·15	0·5596	0·55	0·7088	0·95	0·8289	1·35	0·9115	1·75	0·9599	2·15	0·98422
·16	·5636	·56	·7123	·96	·8315	·36	·9131	·76	·9608	·16	·98461
·17	·5675	·57	·7157	·97	·8340	·37	·9147	·77	·9616	·17	·98500
·18	·5714	·58	·7190	·98	·8365	·38	·9162	·78	·9625	·18	·98537
·19	·5753	·59	·7224	·99	·8389	·39	·9177	·79	·9633	·19	·98574
0·20	0·5793	0·60	0·7257	1·00	0·8413	1·40	0·9192	1·80	0·9641	2·20	0·98610
·21	·5832	·61	·7291	·01	·8438	·41	·9207	·81	·9649	·21	·98645
·22	·5871	·62	·7324	·02	·8461	·42	·9222	·82	·9656	·22	·98679
·23	·5910	·63	·7357	·03	·8485	·43	·9236	·83	·9664	·23	·98713
·24	·5948	·64	·7389	·04	·8508	·44	·9251	·84	·9671	·24	·98745
0·25	0·5987	0·65	0·7422	1·05	0·8531	1·45	0·9265	1·85	0·9678	2·25	0·98778
·26	·6026	·66	·7454	·06	·8554	·46	·9279	·86	·9686	·26	·98809
·27	·6064	·67	·7486	·07	·8577	·47	·9292	·87	·9693	·27	·98840
·28	·6103	·68	·7517	·08	·8599	·48	·9306	·88	·9699	·28	·98870
·29	·6141	·69	·7549	·09	·8621	·49	·9319	·89	·9706	·29	·98899
0·30	0·6179	0·70	0·7580	1·10	0·8643	1·50	0·9332	1·90	0·9713	2·30	0·98928
·31	·6217	·71	·7611	·11	·8665	·51	·9345	·91	·9719	·31	·98956
·32	·6255	·72	·7642	·12	·8686	·52	·9357	·92	·9726	·32	·98983
·33	·6293	·73	·7673	·13	·8708	·53	·9370	·93	·9732	·33	·99010
·34	·6331	·74	·7704	·14	·8729	·54	·9382	·94	·9738	·34	·99036
0·35	0·6368	0·75	0·7734	1·15	0·8749	1·55	0·9394	1·95	0·9744	2·35	0·99061
·36	·6406	·76	·7764	·16	·8770	·56	·9406	·96	·9750	·36	·99086
·37	·6443	·77	·7794	·17	·8790	·57	·9418	·97	·9756	·37	·99111
·38	·6480	·78	·7823	·18	·8810	·58	·9429	·98	·9761	·38	·99134
·39	·6517	·79	·7852	·19	·8830	·59	·9441	·99	·9767	·39	·99158
0·40	0·6554	0·80	0·7881	1·20	0·8849	1·60	0·9452	2·00	0·9772	2·40	0·99180

TABLE 4. THE NORMAL DISTRIBUTION FUNCTION

x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$	x	$\Phi(x)$
2·40	0·99180	2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918
·41	·99202	·56	·99477	·71	·99664	·86	·99788	·01	·99869	·16	·99921
·42	·99224	·57	·99492	·72	·99674	·87	·99795	·02	·99874	·17	·99924
·43	·99245	·58	·99506	·73	·99683	·88	·99801	·03	·99878	·18	·99926
·44	·99266	·59	·99520	·74	·99693	·89	·99807	·04	·99882	·19	·99929
2·45	0·99286	2·60	0·99534	2·75	0·99702	2·90	0·99813	3·05	0·99886	3·20	0·99931
·46	·99305	·61	·99547	·76	·99711	·91	·99819	·06	·99889	·21	·99934
·47	·99324	·62	·99560	·77	·99720	·92	·99825	·07	·99893	·22	·99936
·48	·99343	·63	·99573	·78	·99728	·93	·99831	·08	·99896	·23	·99938
·49	·99361	·64	·99585	·79	·99736	·94	·99836	·09	·99900	·24	·99940
2·50	0·99379	2·65	0·99598	2·80	0·99744	2·95	0·99841	3·10	0·99903	3·25	0·99942
·51	·99396	·66	·99609	·81	·99752	·96	·99846	·11	·99906	·26	·99944
·52	·99413	·67	·99621	·82	·99760	·97	·99851	·12	·99910	·27	·99946
·53	·99430	·68	·99632	·83	·99767	·98	·99856	·13	·99913	·28	·99948
·54	·99446	·69	·99643	·84	·99774	·99	·99861	·14	·99916	·29	·99950
2·55	0·99461	2·70	0·99653	2·85	0·99781	3·00	0·99865	3·15	0·99918	3·30	0·99952

The critical table below gives on the left the range of values of x for which $\Phi(x)$ takes the value on the right, correct to the last figure given; in critical cases, take the upper of the two values of $\Phi(x)$ indicated.

3·075	0·9990	3·263	0·9994	3·731	0·99990	3·916	0·99995
3·105	0·9991	3·320	0·9995	3·759	0·99991	3·976	0·99996
3·138	0·9992	3·389	0·9996	3·791	0·99992	4·055	0·99997
3·174	0·9993	3·480	0·9997	3·826	0·99993	4·173	0·99998
3·215	0·9994	3·615	0·9998	3·867	0·99994	4·417	0·99999
			0·9999		0·99995		1·00000

When $x > 3{\cdot}3$ the formula $1-\Phi(x) \doteqdot \frac{e^{-\frac{1}{2}x^2}}{x\sqrt{2\pi}}\left[1-\frac{1}{x^2}+\frac{3}{x^4}-\frac{15}{x^6}+\frac{105}{x^8}\right]$ is very accurate, with relative error less than $945/x^{10}$.

TABLE 5. PERCENTAGE POINTS OF THE NORMAL DISTRIBUTION

This table gives percentage points $x(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{2\pi}}\int_{x(P)}^{\infty} e^{-\frac{1}{2}t^2}\,dt.$$

If X is a variable, normally distributed with zero mean and unit variance, $P/100$ is the probability that $X \geqslant x(P)$. The lower P per cent points are given by symmetry as $-x(P)$, and the probability that $|X| \geqslant x(P)$ is $2P/100$.



P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$	P	$x(P)$
50	0·0000	5·0	1·6449	3·0	1·8808	2·0	2·0537	1·0	2·3263	0·10	3·0902
45	0·1257	4·8	1·6646	2·9	1·8957	1·9	2·0749	0·9	2·3656	0·09	3·1214
40	0·2533	4·6	1·6849	2·8	1·9110	1·8	2·0969	0·8	2·4089	0·08	3·1559
35	0·3853	4·4	1·7060	2·7	1·9268	1·7	2·1201	0·7	2·4573	0·07	3·1947
30	0·5244	4·2	1·7279	2·6	1·9431	1·6	2·1444	0·6	2·5121	0·06	3·2389
25	0·6745	4·0	1·7507	2·5	1·9600	1·5	2·1701	0·5	2·5758	0·05	3·2905
20	0·8416	3·8	1·7744	2·4	1·9774	1·4	2·1973	0·4	2·6521	0·01	3·7190
15	1·0364	3·6	1·7991	2·3	1·9954	1·3	2·2262	0·3	2·7478	0·005	3·8906
10	1·2816	3·4	1·8250	2·2	2·0141	1·2	2·2571	0·2	2·8782	0·001	4·2649
5	1·6449	3·2	1·8522	2·1	2·0335	1·1	2·2904	0·1	3·0902	0·0005	4·4172

TABLE 10. PERCENTAGE POINTS OF THE t-DISTRIBUTION

This table gives percentage points $t_\nu(P)$ defined by the equation

$$\frac{P}{100} = \frac{1}{\sqrt{\nu\pi}} \frac{\Gamma(\frac{1}{2}\nu+\frac{1}{2})}{\Gamma(\frac{1}{2}\nu)} \int_{t_\nu(P)}^{\infty} \frac{dt}{(1+t^2/\nu)^{\frac{1}{2}(\nu+1)}}.$$

Let X_1 and X_2 be independent random variables having a normal distribution with zero mean and unit variance and a χ^2-distribution with ν degrees of freedom respectively; then $t = X_1/\sqrt{X_2/\nu}$ has Student's t-distribution with ν degrees of freedom, and the probability that $t \geqslant t_\nu(P)$ is $P/100$. The lower percentage points are given by symmetry as $-t_\nu(P)$, and the probability that $|t| \geqslant t_\nu(P)$ is $2P/100$.



The limiting distribution of t as ν tends to infinity is the normal distribution with zero mean and unit variance. When ν is large interpolation in ν should be harmonic.

P	40	30	25	20	15	10	5	2·5	1	0·5	0·1	0·05
$\nu = 1$	0·3249	0·7265	1·0000	1·3764	1·963	3·078	6·314	12·71	31·82	63·66	318·3	636·6
2	0·2887	0·6172	0·8165	1·0607	1·386	1·886	2·920	4·303	6·965	9·925	22·33	31·60
3	0·2767	0·5844	0·7649	0·9785	1·250	1·638	2·353	3·182	4·541	5·841	10·21	12·92
4	0·2707	0·5686	0·7407	0·9410	1·190	1·533	2·132	2·776	3·747	4·604	7·173	8·610
5	0·2672	0·5594	0·7267	0·9195	1·156	1·476	2·015	2·571	3·365	4·032	5·893	6·869
6	0·2648	0·5534	0·7176	0·9057	1·134	1·440	1·943	2·447	3·143	3·707	5·208	5·959
7	0·2632	0·5491	0·7111	0·8960	1·119	1·415	1·895	2·365	2·998	3·499	4·785	5·408
8	0·2619	0·5459	0·7064	0·8889	1·108	1·397	1·860	2·306	2·896	3·355	4·501	5·041
9	0·2610	0·5435	0·7027	0·8834	1·100	1·383	1·833	2·262	2·821	3·250	4·297	4·781
10	0·2602	0·5415	0·6998	0·8791	1·093	1·372	1·812	2·228	2·764	3·169	4·144	4·587
11	0·2596	0·5399	0·6974	0·8755	1·088	1·363	1·796	2·201	2·718	3·106	4·025	4·437
12	0·2590	0·5386	0·6955	0·8726	1·083	1·356	1·782	2·179	2·681	3·055	3·930	4·318
13	0·2586	0·5375	0·6938	0·8702	1·079	1·350	1·771	2·160	2·650	3·012	3·852	4·221
14	0·2582	0·5366	0·6924	0·8681	1·076	1·345	1·761	2·145	2·624	2·977	3·787	4·140
15	0·2579	0·5357	0·6912	0·8662	1·074	1·341	1·753	2·131	2·602	2·947	3·733	4·073
16	0·2576	0·5350	0·6901	0·8647	1·071	1·337	1·746	2·120	2·583	2·921	3·686	4·015
17	0·2573	0·5344	0·6892	0·8633	1·069	1·333	1·740	2·110	2·567	2·898	3·646	3·965
18	0·2571	0·5338	0·6884	0·8620	1·067	1·330	1·734	2·101	2·552	2·878	3·610	3·922
19	0·2569	0·5333	0·6876	0·8610	1·066	1·328	1·729	2·093	2·539	2·861	3·579	3·883
20	0·2567	0·5329	0·6870	0·8600	1·064	1·325	1·725	2·086	2·528	2·845	3·552	3·850
21	0·2566	0·5325	0·6864	0·8591	1·063	1·323	1·721	2·080	2·518	2·831	3·527	3·819
22	0·2564	0·5321	0·6858	0·8583	1·061	1·321	1·717	2·074	2·508	2·819	3·505	3·792
23	0·2563	0·5317	0·6853	0·8575	1·060	1·319	1·714	2·069	2·500	2·807	3·485	3·768
24	0·2562	0·5314	0·6848	0·8569	1·059	1·318	1·711	2·064	2·492	2·797	3·467	3·745
25	0·2561	0·5312	0·6844	0·8562	1·058	1·316	1·708	2·060	2·485	2·787	3·450	3·725
26	0·2560	0·5309	0·6840	0·8557	1·058	1·315	1·706	2·056	2·479	2·779	3·435	3·707
27	0·2559	0·5306	0·6837	0·8551	1·057	1·314	1·703	2·052	2·473	2·771	3·421	3·690
28	0·2558	0·5304	0·6834	0·8546	1·056	1·313	1·701	2·048	2·467	2·763	3·408	3·674
29	0·2557	0·5302	0·6830	0·8542	1·055	1·311	1·699	2·045	2·462	2·756	3·396	3·659
30	0·2556	0·5300	0·6828	0·8538	1·055	1·310	1·697	2·042	2·457	2·750	3·385	3·646
32	0·2555	0·5297	0·6822	0·8530	1·054	1·309	1·694	2·037	2·449	2·738	3·365	3·622
34	0·2553	0·5294	0·6818	0·8523	1·052	1·307	1·691	2·032	2·441	2·728	3·348	3·601
36	0·2552	0·5291	0·6814	0·8517	1·052	1·306	1·688	2·028	2·434	2·719	3·333	3·582
38	0·2551	0·5288	0·6810	0·8512	1·051	1·304	1·686	2·024	2·429	2·712	3·319	3·566
40	0·2550	0·5286	0·6807	0·8507	1·050	1·303	1·684	2·021	2·423	2·704	3·307	3·551
50	0·2547	0·5278	0·6794	0·8489	1·047	1·299	1·676	2·009	2·403	2·678	3·261	3·496
60	0·2545	0·5272	0·6786	0·8477	1·045	1·296	1·671	2·000	2·390	2·660	3·232	3·460
120	0·2539	0·5258	0·6765	0·8446	1·041	1·289	1·658	1·980	2·358	2·617	3·160	3·373
∞	0·2533	0·5244	0·6745	0·8416	1·036	1·282	1·645	1·960	2·326	2·576	3·090	3·291